



factsheet

Coma and reduced awareness states

Introduction

Whether it lasts for a few seconds or a few weeks, the usual immediate effect of brain injury is a loss of consciousness. Coma can be defined as a state of depressed consciousness where a person is unresponsive to the outside world. Consciousness relies on the cerebral hemispheres interacting with an area of the brain stem called the ascending reticular activating system. Injury to these areas causes decreased consciousness or coma.

Levels of coma

There are different levels of coma, ranging from very deep, where the patient shows no response or awareness at all, to shallower levels, where the patient responds to stimulation by movement or opening eyes. Still shallower levels can occur, where the patient is able to make some response to speech. Level of coma is usually initially assessed by the Glasgow Coma Scale (GCS).

The GCS is a very simple, easy to administer technique which is used to rate the severity of coma. It assesses the patient's ability to open their eyes, move and speak. The total score is calculated by adding up the scores from the different categories, shown in the table below, and ranges from a minimum of 3 to a maximum of 15.

Best motor response

6	Obeying commands
5	Movement localised to stimulus
4	Withdraws
3	Abnormal muscle bending and flexing
2	Involuntary muscle straightening and extending
1	None

Verbal responses

5	Orientated response
4	Confused conversation
3	Inappropriate words
2	Incomprehensible sounds
1	None

Eye opening

4	Spontaneous
3	To speech
2	To pain
1	None



Recovery from coma

Recovery from coma is a gradual process, starting with the person's eyes opening, then responding to pain, and then responding to speech. People do not just wake up from a coma, and say, 'Where am I?' as is sometimes portrayed in films. The length of coma is one of the most accurate predictors of the severity of long-term symptoms. The longer the coma, the greater the likelihood of residual symptoms, particularly physical disabilities, although this is only a guide and some people can make good recoveries after an extended period in a coma.

Post-traumatic amnesia

After a coma, during a period known as post-traumatic amnesia (PTA), the patient's behaviour may well be restless, disinhibited and agitated. Uncharacteristic behaviour such as swearing, shouting and inappropriate sexual behaviour are not unusual, but are these best ignored, as seeing other people's distress may only increase the patient's agitation or distress. An individual cannot be held responsible for their behaviour during this period. This is a difficult time for relatives, but it is important to remember that the patient will come out of it.

Length of PTA, as with length of coma, is important. This is the best indicator of severity of brain injury. PTA is assessed by asking the patient a number of questions at regular intervals. The first group of questions is concerned with awareness of time, place and person, for example, 'What is your name?', 'What day of the week is it?' A second group of questions relates to the patient's awareness of the accident, e.g. 'What was your last memory before the accident?' A patient deep in PTA will not be able to answer these questions correctly. As the patient emerges from PTA, the answers will become more accurate and more sensible. For more information on PTA, see the Headway factsheet 'Post-traumatic amnesia'.

Vegetative state

A small number of people sustain a brain injury so severe that, although they emerge from coma and have sleep-wake cycles, they have no conscious awareness of themselves or their surroundings. If this condition persists for more than four weeks they can be classified as being in a persistent vegetative state. If it continues for 12 months after traumatic brain injury or 6 months after non-traumatic brain injury, the person can be classed as being in a permanent vegetative state. If the person shows no signs of recovery at this point and staff and family members agree then it is possible to gain a court order to withdraw treatment.



Minimally conscious state (MCS)

This is a condition distinct from coma and vegetative state as the person shows distinct but limited signs awareness and response to stimulation. However, they find it very difficult to remain aware or responsive for any length of time or in a predictable way. People often enter minimally conscious state after being in a vegetative state.

Some in MCS should be able to do at least one of the following:

- Follow simple commands
- Answer simple 'yes' or 'no' questions using either simple words or gestures
- Communicate in some intelligible way
- Show some purposeful actions, such as reaching for an object

Locked-in syndrome

This is a rare condition in which a person is aware of themselves and their surroundings but is unable to move or speak. Often people can move their eyes or eyelids and may be able to communicate by blinking.

Conclusion

Having a relative in a coma, or some other form of reduced awareness state, is a very distressing and confusing time. It is very important to communicate with the medical staff and to understand as much as possible about the person's level of awareness. However, the most important thing is to look after yourself and seek help and support.

Vegetative states, minimally conscious states and locked-in syndrome are thankfully relatively rare. These conditions are not fully understood and accurate diagnosis can sometimes be difficult. For example, some people who have been thought to be in a vegetative state have subsequently been shown to possess some awareness. However, that does not mean that everyone does. Although most families cling to the idea that there is some awareness, lack of awareness could be preferable for a time. There is often hope of improvement in the future.

If you would like to discuss the issues raised in this factsheet further, please contact the Headway helpline on **0808 800 2244** or helpline@headway.org.uk.



factsheet

Further reading

The following books are available from Headway and provide a good introduction to brain injury and its effects, including sections on coma.

Daisley, A., Tams, R. and Kischka, U. (2008) *Head Injury: The Facts*. Oxford: Oxford University Press.

Powell, T. (2004) *Head Injury: A Practical Guide*. Milton Keynes: Speechmark Publishing Ltd.

Headway also produces an extensive range of booklets and factsheets covering the range of problems that brain injury can cause. Booklets that may be of particular interest include:

- *Caring for someone with a brain injury*
- *Coma stimulation activities*
- *Effects of Brain Injury and How to Help*
- *Hospital Treatment and Early Recovery after Brain Injury*
- *Rehabilitation after Brain Injury*

To order copies of books and booklets please visit our website at www.headway.org.uk/shop.aspx, or telephone 0115 924 0800.

Factsheets are free to download at www.headway.org.uk/factsheets.aspx and copies can also be obtained from the Headway helpline on 0808 800 2244 or helpline@headway.org.uk.



About Headway

Headway - the brain injury association is a charity set up to give help and support to people affected by brain injury.

A network of local Headway Groups and Branches throughout the UK offers a wide range of services including rehabilitation programmes, carer support, social re-integration, community outreach and respite care.

The Headway helpline provides information, advises on sources of support, finds local rehabilitation services and offers a listening ear to those experiencing problems.

Other services provided by Headway include:

- Supporting and developing local Groups and Branches
- Promoting understanding of brain injury and its effects
- Producing a range of publications on aspects of brain injury
- Lobbying for better support and resources to be made available by statutory health and social care providers
- Campaigning for measures that will reduce the incidence of brain injury
- Accreditation of UK care providers through the Approved Provider scheme

Make a donation

I wish to help Headway - the brain injury association to improve life after brain injury by making a donation £

Your personal details

Mr/Mrs/Miss/Ms

Full name

Address

Postcode

Daytime tel. number

Email

Your payment details

I enclose my cheque for £ made payable to Headway - the brain injury association

Please debit my Visa / Mastercard / Diners / Maestro card number:

Security code*:

*3 digits on reverse
of card

Expiry date: / Start date: / Issue no. (Maestro only):

We (Headway - the brain injury association) hold the information you supply to us on our database. From time to time we may send you information regarding other services and products we provide. If you do not wish to receive such information, please tick this box.

Please tick this box if you do not want a receipt

Gift Aid

gift aid it Gift Aid will increase the value of your donation by 25% at no extra cost to you. Please tick the Gift Aid box below if you want Headway to reclaim the tax you have paid on your donations since 6th April 2008 and any future donations you make. Please note that you must pay an amount of income tax or capital gains tax at least equal to the amount we can reclaim on your donation in any tax year.

I confirm I have paid or will pay an amount of income tax and/or capital gains tax for the current tax year (6 April to 5 April) that is at least equal to the amount of tax that all the charities and Community Amateur Sports Clubs (CASCs) that I donate to will reclaim on my gifts for the current tax year. I understand that other taxes such as VAT and council tax do not qualify. I understand the charity will reclaim 25p of tax on every £1 that I have given.

Donate by phone or online

- To make a donation by phone, call:
0115 924 0800
- To make a donation online visit:
www.headway.org.uk

Completed forms should be sent to: The Fundraising Team, Headway - the brain injury association, FREEPOST RSGU-CSKS-CSJS, Bradbury House, 190 Bagnall Road, Old Basford, Nottingham, NG6 8SF